

Name _____
PHYS 0111 (Gen Chem II)

Test 2
Spring 2005

A few equations:

1. _____

$$K = \frac{[\text{products}]}{[\text{reactants}]}$$

2. _____

$$t_{1/2} = \frac{\ln(2)}{k}$$

3. _____

$$t_{1/2} = [A]_0 / (2k)$$

4. _____

$$t_{1/2} = \frac{1}{k[A]_0}$$

5. _____

$$\frac{\text{rate}_1}{\text{rate}_2} = \frac{[\text{conc}_1]^x}{[\text{conc}_2]^x}$$

6. _____

Integral Rate Laws

$$\ln[A]_t = -kt + \ln[A]_0$$

$$1/[A]_t = kt + 1/[A]_0$$


$$[A]_t = -kt + [A]_0$$

Differential Rate Laws (general form)


$$\text{rate} = k[A]^x[B]^y$$

1. (5 pts. ea.) Write the equilibrium expression, K, for the following reactions.

a.




b.



c.



2. Rate and concentration data were collected for the following reaction.



Experiment	Initial [NO]	Initial [H ₂]	Initial -d[NO]/dt (M/s)
1	0.24	0.10	14.4 x 10 ⁻⁶
2	0.12	0.10	3.6 x 10 ⁻⁶
3	0.12	0.15	5.4 x 10 ⁻⁶


a. (3 pts.) For experiment 1, determine the rate at which N₂ is being produced.

b. (10 pts.) Determine the differential rate law for the reaction. Do not determine the rate constant.

3. The concentration of NOBr was measured as the following reaction proceeded, and the data is tabulated below.




Time (s)	0	10	20	30	40
[NOBr]	0.0400	0.0303	0.0244	0.0204	0.0175




$$y = -0.0005x + 0.0375$$

$$R^2 = 0.9439$$



$$y = -0.0205x - 3.2635$$

$$R^2 = 0.9859$$



$$y = 0.803x + 24.969$$


$$R^2 = 0.9999$$

a. (6 pts.) What is the order of the reaction with respect to NOBr concentration?

b. (4 pts.) How long does it take for the NOBr concentration to reach 0.0105 M?

c. (4 pts.) Determine the differential rate law. Include the value for the rate constant with the appropriate units.

4. In tetrahydrofuran (THF), an organic solvent, $\text{Mo}(\text{CO})_6$ reacts with $\text{P}(\text{CH}_3)_3$ to form $\text{Mo}(\text{CO})_5(\text{P}(\text{CH}_3)_3)$. Experiments reveal that the rate of the reaction is 0 order with respect to the $\text{P}(\text{CH}_3)_3$ concentration and is 1st order with respect to $\text{Mo}(\text{CO})_6$ concentration.




Two mechanisms have been proposed for the reaction.

Mechanism I



Mechanism II



- a. (3 pts) Write the experimental rate law.
- b. (4 pts.) Determine the rate law predicted by mechanism I .
- c. (4 pts.) Determine the rate law predicted by mechanism II.
- d. (3 pts.) Which mechanism is incorrect?

5. (2 pts ea.) Identify whether each of the following statements about a dynamic equilibrium is true or false.

_____ At equilibrium, the concentrations of the reactants and the products do not change.

_____ At equilibrium, the rate at which the reactants are converted to products is the same as the rate at which products are converted to reactants.

_____ At equilibrium, the rate constant for the forward reaction is always equal to the rate constant for the reverse reaction.

_____ At equilibrium, the rate at which the reactants are converted to products is the same as the rate at which products are converted to reactants.

_____ At equilibrium, the forward and reverse reactions stop.

_____ A large equilibrium constant means that when equilibrium is reached there will be more reactants than products present.

_____ A small equilibrium constant means that when equilibrium is reached there will be more reactants than products present.

6. (2 pts. ea.) The following statements refer to the relationship between a reaction's activation energy, E_a , the rate constant for the reactions, k , and temperature, T . Identify whether each statement is true or false.

_____ In general, reactions with large activation energies are slow.

_____ A reaction with a large rate constant is an intrinsically fast reaction.

_____ Typically, reactions with large rate constants have small activation energies.

_____ In general, as the temperature of a reaction is increased, the rate of the reaction is increased.

_____ As the temperature of a reaction is increased, the activation energy barrier is lowered.

_____ As the temperature of a reaction is increased, more reactant molecules are capable of overcoming the activation energy barrier.

_____ In general, increasing the temperature of a reaction increases the reaction's rate constant.